

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

AB

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/85, C07K 16/12, 16/46, C12N 5/10, A61K 39/40		A1	(11) International Publication Number: WO 99/32645 (43) International Publication Date: 1 July 1999 (01.07.99)
(21) International Application Number: PCI/US98/27267 (22) International Filing Date: 22 December 1998 (22.12.98) (30) Priority Data: 60/068,635 23 December 1997 (23.12.97) US 09/215,163 18 December 1998 (18.12.98) US (71)(72) Applicants and Inventors: STINSON, Jeffrey, L. [US/US]; 15030 Durham Lane, Davie, FL 33331 (US). WONG, Hing [US/US]; 2966 Wentworth, Weston, FL 33332 (US). O'BRIEN, Alison, D. [US/US]; 5514 Charlcote Road, Bethesda, MD 20817 (US). SCHMITT, Clare, K. [US/US]; 4616 Glasgow Drive, Rockville, MD 20853 (US). MELTON-CELSA, Angela [US/US]; 20516 Brandywine Court, Sterling, VA 20165 (US). (74) Agents: GARRETT, Arthur, S. et al. ; Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P., 1300 I Street, N.W., Washington, DC 20005-3315 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: HUMANIZED MONOCLONAL ANTIBODIES THAT PROTECT AGAINST SHIGA TOXIN INDUCED DISEASE			
(57) Abstract <p>The present invention describes the preparation and use of biologically and immunologically active humanized monoclonal antibodies to Shiga toxin, a toxin associated with HC and the potentially life-threatening sequela HUS transmitted by strains of pathogenic bacteria. The present invention describes how these humanized antibodies may be used in the treatment or prevention of Shiga toxin induced diseases. One aspect of the invention is the humanized monoclonal antibody which binds Shiga toxin where the constant regions are IgG1-kappa and the variable regions are murine in origin. Yet another aspect of the invention is expression vectors and host cells transformed with such vectors which express the humanized monoclonal antibodies of the present invention.</p>			
<p>13C4 ANTI-STX1 HEAVY CHAIN VARIABLE REGION</p> <p>CAGGTGCAGCTGCAGGAGTCTGGGGCTGAGCTGGAGTCTGGGGCTCAGTGAGGATC SEQ ID NO.18 Q V Q L Q E S G A E L V R S G A S V R H SEQ ID NO.19 TCCTGCAGGCTTCTGGCTACACATTTACCACTTACAATATGCACTGGGTAAACAGACA S C D A S G Y T F T S Y N M H W V K Q T CCTGGACAGGCTTGAATGGATGATATATTTATCTGGAAATGGTGGTACTAACTAC P G Q G L E W I G Y I Y P G N G G T N Y ATTGAGAAATTTAAGGCAAGCCATATGACTGCAGACATCTCTCCAGCACAGCTAC I Q K F K G K A I L T A D T S S S T A Y ATGCAAGTACAGCTCTGACATCTGAAGACTCTCGGCTATTTCTGTACAAGAGTCCCTCT M Q I S S L T S E D S A V Y F C T R S P S CACTACAGTAGTGACCCCTACTTTGACTACTGGGCCAGGCAACACTCTCAGCTCTCTCA H Y S S D P Y F D Y W G Q G T T L T V S S</p> <p>13C4 ANTI-STX1 LIGHT CHAIN VARIABLE REGION</p> <p>GATATCGTGATGCCAGCTCTCAAAATTCATGTCACATCAGTCGGAGACAGGCTCAGCATC SEQ ID NO.20 D I V M S Q S H K F M S T S V G D R V S I SEQ ID NO.21 ACCTGTAAGGCCAGCCAGATGTGGTACTGCTGTGCTGATCAGCAGAAATCCAGGACAA T C K A S Q D V G T A V A W Y Q N P G Q TCTCTAAATTTCTGATTTACTGGGCATCCACAGGCACACTGGAGTCCCTGATCCCTTACA S P K F L I Y W A S T R H T G V P D R P T GGCAGTGGATCTGGGACAGATTCTACTCTCACCATTACCAATGTGCACTCTGAAGACTTGGCA G S G S G T D F T L T I T N V Q S E D L A GATTATTTCTGTGAGCAATATAGCAGTTATCTCTACGTTGGTGGTGGGACCACTCTGGAG D Y P C Q Q Y S S Y P L T F G A G T S L E CTGAACGT L K R</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HUMANIZED MONOCLONAL ANTIBODIES THAT PROTECT AGAINST SHIGA TOXIN INDUCED DISEASE

Related Application

This application claims the benefit of U.S. Provisional Application No. 60/068,635, filed December 23, 1997, which is hereby incorporated by reference.

Government Interest

5 The invention described herein may be manufactured, licensed, and used for governmental purposes without payment of royalties to us thereon.

Field of the Invention

The invention relates to a family of multi-unit bacterial proteins that are associated with hemorrhagic colitis and the life-threatening sequela, hemolytic uremic syndrome. These
10 proteins, defined as members of the "Shiga toxin family", have been utilized in the isolation and identification of murine monoclonal antibodies. The invention further relates to the construction of humanized monoclonal antibodies which incorporate the mouse variable regions. It also relates to antibodies to Shiga toxins or toxoids, both monoclonal and polyclonal, and their use in treating, diagnosing, and preventing of disease and infections caused by pathogenic *E. coli*.
15 Finally, the invention relates to preparing the humanized monoclonal antibodies to proteins of the Shiga toxin family.

Background of the Invention

Enterohemorrhagic *Escherichia coli* (EHEC) are associated with food-borne outbreaks of bloody diarrhea or "hemorrhagic colitis" (HC) and the hemolytic uremic syndrome (HUS).
20 (Spika, J. et al., *J. Pediatr.*, 109: 287-291(1986); Remis, R., *Ann. Intern. Med.*, 101:624-626

(1984); Riley, L. et al., *N. Engl. J. Med.*, 308:681-685 (1983)). EHEC infection can be deadly and poses a significant threat to the young and the elderly, who are the most likely to develop serious complications from EHEC infections. Several outbreaks and sporadic cases of HC and HUS have occurred over the past few years, with the largest U.S. outbreak in 1993. In that
5 outbreak, over 500 cases of HC and HUS were traced to contaminated hamburgers from a Jack-in-the Box fast food restaurant. (Centers for Disease Control and Prevention, *Morbid. Mortal. Weekly Rep.*, 42:258(1993)). In July 1996, a large outbreak of EHEC in Japan resulted in over 10,000 infected individuals and 8 deaths. Many Japanese children required hospitalization.

Primarily, HC and HUS are transmitted by the ingestion of contaminated food,
10 particularly undercooked beef products, such as hamburger. (Doyle et al., *J. Appl. Environ. Microbiol.* 53:2394 (1987); Samadpour et al., *J. Appl. Environ. Microbiol.* 60:1038 (1994)). With the prevalence of EHEC in cattle and the subjective nature of differentiating between cooked and undercooked hamburgers, a stop at a fast food restaurant or a family barbecue can result in tragedy.

15 HC and HUS appear to be mediated by the toxin produced by EHEC and *Shigella dysenteriae* (for a review, see O'Brien and Holmes, *Microbiol. Rev.*, 51:206-220 (1987)). These bacteria produce a family of closely related cytotoxins that collectively will be called "Shiga toxins" for the purpose of this application. Shiga toxins (alternatively, "verotoxins") have cytotoxic, and enterotoxic activity (Strockbine, N. et al., *Infect. Immun.*, 53:135-140 (1986)).

20 Based on the exhibition of immunological non-cross-reactivity, the Shiga toxins have been divided into two groups. (Strockbine et al., *supra*). These groups have been designated Shiga toxin type 1 (Stx1) and Shiga toxin type 2 (Stx2). The Stx1 group contains the prototype Stx1 toxin from EHEC as well as the Shiga toxin from *Shigella dysenteriae* type 1. In recent years, other types of toxins have been discovered and considered to be members of the Stx2

group. These are Stx2e, Stx2c, and Stx2d. (Lindgren et al., *Infection and Immunology*, 61:3832 (1993); Schmitt, C. et al., *Infect. Immun.*, 59:1065-1073 (1991); Marques, L. et al., *FEMS Lett.*, 44:33-38 (1987)).

For the purposes of this application the term "Shiga toxin" encompasses Shiga toxin and
5 any other toxins in the Stx1 or Stx2 group or their variants. The abbreviation "Stx" will refer to the toxin protein itself.

Despite this knowledge about the results of exposure to these toxins, currently there is no known cure or vaccine for HC or HUS. Antibiotics may even make the severe complications worse by increasing toxin release from bacteria. Thus, there is a need for a compound to prevent
10 or to treat the complications of EHEC produced by Shiga toxin. Such a compound could be used to treat infected patients and decrease the systemic effects of toxin on the CNS, blood and kidneys. In addition, if the toxin could be neutralized, antibiotics could be safely given to kill the bacteria in the GI tract. Such a compound could also be used to prevent infectious complications, by treating exposed or high risk individuals before they acquire EHEC infection.
15 Such individuals would include children in day care or the elderly in nursing homes, where a case of EHEC diarrhea has been detected. These individuals are at increased risk to develop severe complications and spread of EHEC in these environments is not unusual.

The knowledge of the immunological cross reactivity of the Shiga toxins offers a tantalizing prospect for pharmacological approaches to EHEC treatment. Currently, there are no
20 known prophylactic or therapeutic agents available for this disease. Accordingly, there is a need in the art to provide monoclonal antibodies that can bind to Shiga toxins which could prevent or lessen the devastating effects of these toxins. There is also a need in the art for data concerning the binding site of such antibodies so that other antibodies with similar abilities can be identified and isolated.

There is a related need in the art for humanized or other chimeric human/mouse monoclonal antibodies. In well publicized studies, patients administered murine anti-TNF (tumor necrosis factor) monoclonal antibodies developed anti-murine antibody responses to the administered antibody. (See, e.g., Exley A.R., et al., *Lancet* 335:1275-1277 (1990)). This type of immune response to the treatment regimen, commonly referred to as the HAMA response (for human anti-mouse antibodies), decreases the effectiveness of the treatment and may even render the treatment completely ineffective. Humanized or chimeric human/mouse monoclonal antibodies have been shown to significantly decrease the HAMA response and to increase the therapeutic effectiveness. (LoBuglio et al., *P.N.A.S.* 86:4220-4224 (June 1989)).

10

Summary of the Invention

To satisfy these needs in the art, applicants provide humanized mouse monoclonal antibodies to Shiga toxin 1 and Shiga toxin 2. In addition, this invention sets forth the variable region associated with Shiga toxin neutralization as well as the specific complementarity determining regions (CDRs) of the variable region. The present invention particularly provides human monoclonal antibodies, derived from the mouse monoclonal antibodies 13C4 and 11E10, which are specifically reactive with Stx1 and Stx2, respectively.

15

In another aspect, the invention includes a pharmaceutical composition and a method of treatment with such composition which will prevent or treat Shiga toxin induced disease.

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory and are intended to provide further explanation of the invention as claimed.

20

The accompanying drawings are included to provide a further understanding of the invention and are incorporated in and constitute a part of this specification. They illustrate

several embodiments of the invention and, together with the description, serve to explain the principles of the invention.

Brief Description of the Drawings

Figure 1 provides a schematic representation of the general cloning strategy used for
5 cloning the variable region gene fragments of both the 11E10 and 13C4 antibodies.

Figure 2 lists the oligonucleotide sequences used for PCR amplification of the variable regions of the 13C4 antibody.

Figure 3 recites the DNA and amino acid consensus sequences for the variable regions of both the heavy and light chains of the 13C4 antibody. The DNA consensus sequence was
10 determined using automated fluorescent dye-terminator sequence reactions of the amplified cDNAs analyzed with an ABI 373 sequencing instrument. The CDRs were identified according to the nomenclature of Kabat and Wu (Kabat, et al. (1991), *Sequences of Proteins of Immunological Interest*, Vol. I, 5th edition, U.S. Department of Health and Human Services) and are located as follows:

15	Heavy Chain CDRs: (SEQ ID NO:19)	CDR1-aa31-35 CDR2-aa50-66 CDR3-aa99-111
20	Light Chain CDRs: (SEQ ID NO:21)	CDR1-aa24-34 CDR2-aa50-56 CDR3-aa89-97

Figure 4 illustrates the strategy for construction of the mammalian expression vector tKMC249A containing the variable regions of the 13C4 antibody.

Figure 5 lists the oligonucleotides used for PCR amplification of the variable regions of the 11E10 antibody.

Figure 6 recites the DNA and amino acid consensus sequences for the variable regions of both the heavy and light chains of the 11E10 antibody. The DNA consensus sequence was determined using automated fluorescent dye-terminator sequence reactions of the amplified cDNAs analyzed with an ABI 373 sequencing instrument. The CDRs were identified according to the nomenclature of Kabat and Wu (Kabat and Wu; *supra*) and are located as follows:

	Heavy Chain CDRs: (SEQ ID NO:44)	CDR1-aa31-35 CDR2-aa50-66 CDR3-aa99-108
10	Light Chain CDRs: (SEQ ID NO:42)	CDR1-aa24-40 CDR2-aa56-62 CDR3-aa95-103

Figure 7 illustrates the strategy for construction of the mammalian expression vector pACE4 containing the variable regions of the 11E10 antibody.

Figure 8 shows the anti-Stx1 13C4 antibody production ELISA and demonstrates that the H13C4 cell line produces an antibody consisting of human IgG and kappa constant domains.

Figure 9 shows the anti-Stx1 13C4 toxin binding activity ELISA and demonstrates that the antibody produced by the H13C4 cell line binds to Stx1 toxin.

Figure 10 shows the anti-Stx2 11E10 antibody production ELISA and demonstrates that the H11E10 cell line produces an antibody consisting of human IgG and kappa constant domains.

Figure 11 shows the anti-Stx2 11E10 toxin binding activity ELISA and demonstrates that the antibody produced by the H11E10 cell line binds to Stx2 toxin.

Figure 12 illustrates the strategy for construction of the plasmid pCDNA3mut.LCPL.LCVK.

Figure 13 illustrates the strategy for construction of the plasmid pCDNA3mut.HCPL.HCV2b.

Figure 14 illustrates the strategy for construction of the plasmids pSUN9 and pSUN9:kappa.

Figure 15 illustrates the strategy for construction of the plasmids pSUN10 and pSUN10:IgG1.

5 Figure 16 illustrates the strategy for construction of the plasmid pJRS311.

Figure 17 illustrates the strategy for construction of the plasmid pJRS315.

Additional features and advantages of the invention will be set forth in the description which follows. The objectives and other advantages of the invention will be realized and attained by the composition of matter and process particularly pointed out in the written
10 description and the claims hereof as well as the appended drawings. To achieve these and other advantages and in accordance with the purpose of the invention, as embodied and broadly described, the invention describes monoclonal humanized antibodies which bind Shiga toxin proteins.

Detailed Description of the Invention

15 The present invention relates to humanized monoclonal antibodies that bind to Shiga toxin proteins, and the use of such antibodies in the treatment or prevention of Shiga toxin-induced diseases. Antibodies are proteinaceous structures made up of two heavy and two light chains. The five different classes of higher vertebrate antibodies—IgM, IgD, IgG, IgA and IgE—are distinguished by their heavy chains (mu, delta, gamma, alpha and epsilon,
20 respectively). Each class has further subclasses; IgG, for example, may be IgG1, IgG2, IgG3, or IgG4, where the heavy chain is gamma 1, gamma 2, gamma 3 or gamma 4, respectively. The pair of light chains in these classes or subclasses may be either kappa or lambda.

Antibodies are further divided into a constant region and a variable region. For both the heavy and light chains, the carboxy-terminal ends make up the constant sequence region, while the amino terminal ends contain the variable sequence region. Within these variable regions, the complementarity determining regions (CDRs) are located which are primarily responsible for the observed antigen binding which is characteristic of antibodies.

"Humanized" monoclonal antibodies mean monoclonal antibodies originally from a non-human source to which human components have been substituted. In a preferred embodiment, the humanized monoclonal antibodies of this invention comprise variable regions which derive from non-human sources and constant regions which derive from human sources.

As set forth above, Shiga toxin proteins (Stx) refer to the family of multi-unit bacterial proteins produced by EHEC and *Shigella dysenteriae* which are associated with outbreaks of Shiga toxin induced diseases. Shiga toxin proteins are meant to encompass Stx of *Shigella dysenteriae*-type 1 and Stx type 1 and type 2, and type 2 variant toxins of *E. coli*.

Shiga toxin-induced diseases of humans include bloody diarrhea, hemorrhagic colitis, hemolytic uremic syndrome, and thrombotic thrombocytopenic purpura.

In a more specific embodiment, the invention relates to humanized monoclonal antibodies that have the same binding specificity of at least two well characterized murine monoclonal antibodies. These two monoclonal antibodies were developed in Dr. Alison O'Brien's laboratory as set forth in Strockbine, N.A. et al., *Infection and Immunity*, 1985, 50:695-700 and Perera, L.P. et al., *J. Clinical Microbiol.*, 1988, 26:2127-2131, and have been deposited at the ATCC as set forth below.

By "same binding specificity" is meant a level of binding sufficiently detectable in a standard binding assay to distinguish between toxin binding and non-specific background

binding as exemplified by appropriate controls (for examples, see Figure 9 and Figure 11).

Those of ordinary skill in the art can readily test for binding levels using routine skills and techniques.

In another preferred embodiment, the humanized antibodies are characterized by their structural features. In one aspect of this embodiment, the constant region is a human constant region and the variable region derives from a rodent, preferably a mouse. Although not limited to these, the four variable regions set forth in the specification, particularly in Figure 3 and Figure 6, are preferred. Of course, the invention includes modifications (i.e., deletions, additions, and substitutions) that do not appreciably diminish the characteristic binding associated with the exemplified variable regions.

In yet another embodiment, the CDRs of the variable region are employed in the invention. As noted above, CDRs are located in the variable regions of both the light and heavy chains and are responsible for antigen binding. In view of the ready ability of those in this art to determine the CDR regions, the present invention is not limited to the CDRs specifically set forth. Indeed, the vectors created by the applicants and described in great detail below are sufficient for use with any CDRs of non-human antibodies to Shiga toxins.

In the most preferred embodiment of the present invention, the humanized monoclonal antibody derives its constant regions from IgG1-kappa and its variable regions from all or part of the sequences as set forth in Figures 3 and 6. In further embodiments, the invention also relates to the expression vectors which code for these monoclonal humanized antibodies, and to host cells which have been transformed with such expression vectors.

Finally, the invention comprises the pharmaceutical compositions and the methods of use of these humanized monoclonal antibodies to treat or prevent Shiga toxin-producing bacterial infections. As a pharmaceutical, this invention includes but is not limited to diluents

and carriers known in the art, such as saline and sucrose solutions appropriate for application to patients. As used herein, "patients" refers to any susceptible mammal, such as dogs, horses, mice, etc., but is particularly preferred to mean humans.

The invention further involves the administration of a therapeutically effective amount, as well as a prophylactically effective amount, of the humanized monoclonal antibody of the invention. As those in the art would recognize, a therapeutically effective amount is a dose that ameliorates edema, thrombocytopenia, and uremia associated with EHEC-mediated HUS. Similarly, a prophylactically effective amount is a dose that prevents exposed individuals from developing these symptoms.

The following examples illustrate specific embodiments of the invention.

Humanization of the Anti-Stx1 antibody 13C4

The following examples 1-3 relate to anti-Stx1 antibody 13C4 and its humanized counterpart H13C4.

EXAMPLE 1. Cloning of the 13C4 variable region cDNAs.

The hybridoma cell producing the "13C4" antibody (Anti-Stx1) was deposited on December 2, 1987, at the American Type Culture Collection, Rockville, MD under Accession No. CRL 1794, and can be obtained from the ATCC, or, as here, from Dr. Alison O'Brien (for details of hybridoma preparation, see Strockbine, N.A. et al., *Infection and Immunity*, 50:695-700 (1985)). A vial of cells was thawed and resuspended in IMDM (Mediatech) complete media supplemented with 10% FBS (Irvine).

Total RNA was isolated from 1×10^7 "13C4" cells using the Midi RNA Isolation kit (Qiagen) following the manufacturer's procedure. The RNA was dissolved in 10 mM Tris,

0.1 mM EDTA (pH 8.4) containing 0.03 U/ μ g Prime RNase Inhibitor (5'-3') to a final concentration of 0.25 μ g/ μ l.

Figure 1 shows the strategy for cloning the variable region gene fragments and Figure 2 lists the oligonucleotide primers used. The "13C4" total RNA (2 μ g) was converted to cDNA by using Superscript II-MMLV Reverse Transcriptase (Life Technologies) and mouse kappa (oKA57, SEQ ID NO:57) and mouse CH1 (JS300, SEQ ID NO: 6) specific priming, according to the manufacturer's procedures. The first strand cDNA synthesis products were then purified using a membrane concentrator device (Amicon Centricon 30 or Millipore UltraFree 15). Of the cDNA recovered, 3 μ l was used as template DNA for PCR. Typical PCR amplification reactions (100 μ l) contained template DNA, 50 pmoles of the appropriate primers (JSS9, JSS10, JSS11, JSS12, JS153 & JSS154—SEQ ID NO:9-12 for light chains; JSS1, JSS2, JSS3, JSS4, JSS8 and oKA-143—SEQ ID NO:1-5 and SEQ ID NO:15 for heavy chains), 2.5 units of ExTaq polymerase (PanVera), 1x ExTaq reaction buffer, 200 μ M dNTP, and 2mM $MgCl_2$. Each of the templates was denatured by an initial incubation at 96°C for 1 min. The heavy chain products were amplified by 40 or 50 thermal cycles of 59 to 72°C for 30 sec., 72°C for 30 sec., then 96°C for 1 min. and a final extension step at 72°C for 5 min. The light chain products were amplified by 6 thermal cycles of 46, 48 or 54°C for 30 sec., 72°C for 30 sec., then 96°C for 1 min followed by 35 step cycles of 60°C for 1 min, then 96°C for 1 min. and a final extension step at 72°C for 5 min. The PCR products from the successful reactions were purified using the Wizard PCR Purification system (Promega) as per the manufacturer's procedure.

The heavy chain and light chain PCR products (approximately 400 bp) were then cloned into a bacterial vector for DNA sequence determination. Ligations of the PCR fragments were carried out into the pCR2.1 T/A style cloning vector (Invitrogen) following

the manufacturer's procedures using 1:1, 3:1 and 5:1 insert to vector molar ratios. One half of each of the ligation reactions was used to transform competent XL1 Blue cells.

Heavy chain plasmid clones containing DNA inserts were identified using diagnostic restriction enzyme digests with EcoRI (New England Biolabs). The DNA sequence of
5 plasmids (tKMC217B) containing inserts of the appropriate size (400 bp) were then determined. The final consensus DNA sequence of the heavy chain variable regions is shown in Figure 3.

The light chain PCR products were identified differently. The hybridoma cell line that expresses the "13C4" antibody was made by fusing mouse splenocytes with the SP2/0
10 myeloma cell line. The SP2/0 cell line transcribes a pseudogene for the kappa light chain. The pseudogene transcript, when converted to cDNA by RT-PCR, contains an AflIII restriction site. Light chain candidate clones (tKMC226 & 227) were digested with AflIII (New England Biolabs) using the manufacturer's procedures to identify clones containing inserts of the appropriate size (403 bp; no AflIII site is present in these products). The final
15 consensus DNA sequence of the light chain variable regions is shown in Figure 3 with the CDRs indicated by underlining.

EXAMPLE 2. Construction of the expression vector tKMC249A.

The heavy and light chain variable regions were then subcloned into mammalian expression plasmid vectors for the production of recombinant chimeric mouse/human
20 antibody molecules. The vectors result in the production of recombinant antibody molecules under the control of CMV transcriptional promoters. The heavy chain molecules are direct cDNA constructs that fuse the variable region sequence directly into the human IgG1 constant domain. The light chain molecules, on the other hand, have a mouse kappa intron region 3' of

the variable region coding fragment. After splicing, the variable region becomes fused to a human kappa constant region exon (Figure 4). The selectable marker for the vector in mammalian cells is aminoglycoside phosphotransferase (neo), using the drug G418 (CellTech).

5 A. Creation of the expression vectors

To create the heavy and light chain expression vectors required DNA fragment ligations and site directed mutagenesis steps. The result was vectors that express both antibody chains with CMV promoter driven transcription. Neomycin resistance serves as a dominant selectable marker for transfection of mammalian cells. In addition, these vectors
10 have been designed to allow convenient cloning of any light chain variable region as EcoRV/BstBI fragment, any heavy chain variable region as a NruI/EcoRI fragment, and any heavy chain constant domain as an EcoRI/NotI fragment. These restriction sites were chosen because they occur rarely (if ever) in human and mouse variable regions. There is a mouse J region/kappa intron fragment fused to a human kappa exon so that after post-transcriptional
15 splicing a mouse human chimeric kappa light chain is produced. Lastly, the vectors were designed to facilitate excision (BglII/NheI) of a whole antibody expression cassette from one vector to be ligated into a second vector cut with BamHI/NheI, creating an expression vector with the apparatus for both chains.

The backbone of the vector was the plasmid pCDNA3 (Invitrogen). This plasmid was
20 cut with HindIII/XhoI and a "light chain polylinker" DNA fragment was inserted to create the starting "light chain vector" pCDNA3.LCPL (see Figure 12). This linker contained the restriction sites HindIII, KpnI, ClaI, PmlI, EcoRV, XmaI, BamHI and XhoI to facilitate subsequent cloning steps. A SmaII/BclI DNA fragment containing a light chain leader, mouse anti-CKMB kappa light chain genomic fragment, and 3' UTR was cloned into the

EcoRV/BamHI sites of pCDNA3.LCPL. The mouse kappa intron, exon, and 3' UTR in this fragment was derived from LCPXK2 received from Dr. Richard Near (Near, R.I. et al., 1990, *Mol. Immunol.* 27:901-909). Mutagenesis was then performed to eliminate an NruI (209), MluI (229), and BstBI (2962) and to introduce an NheI (1229) and a BamHI (1214) site to
5 create the plasmid pCDNA3mut.LCPL.LCVK (see Figure 12).

A second "heavy chain vector" pCDNA3mut.HCPL was constructed from the pCDNA3mut.LCPL.LCVK plasmid by replacing the light chain expression region (HindIII/XhoI) with a "heavy chain polylinker" consisting of restriction sites HpaI, BspEI, EcoRV, KpnI, and XhoI. This plasmid was digested with EcoRV and KpnI. A SmaI/KpnI
10 digested DNA fragment containing a heavy chain leader and an anti-CKMB IgG2b mouse heavy chain genomic fragment was then ligated into the EcoRV/KpnI digested plasmid. A KpnI/SalI oligonucleotide fragment containing a 3' UTR and a NotI upstream of the SalI site was subsequently cloned into the KpnI/XhoI digested plasmid, (knocking out the XhoI site), to create the plasmid pCDNA3mut.HCPL.HCV2b (see Figure 13).

15 A human kappa light chain constant domain was then cloned into pCDNA3mut.LCPL.LCVK as a EcoNI/XhoI fragment generating the plasmid tKMC180C2. A human IgG1 constant domain was cloned into pSUN10 as a EcoRI/NotI fragment creating the plasmid pJRS313. The variable regions of 13C4 were cloned into these two vectors (as described above). A BglII/NheI fragment from the human heavy chain vector tKMC229C
20 was then cloned into the human light chain vector tKMC231D cut BamHI/NheI to create tKMC249A (see Figure 4).

B. Subcloning the variable regions into the expression vectors

The variable region gene fragments were re-amplified by PCR using primers that adapted the fragments for cloning into the expression vectors (see Figures 2 and 4). The

heavy chain front primer (oKA143, SEQ ID NO:15) includes a 5' tail that encodes the C-terminus of the heavy chain leader and an NruI restriction site for cloning, while the heavy chain reverse primer (oKA144, SEQ ID NO:14) adds a 3' EcoRI restriction site for cloning. The light chain front primer (oKA145, SEQ ID NO:16) introduces an EcoRV restriction site at the N-terminus of the light chain variable region for cloning, while the light chain reverse primer (oKA146, SEQ ID NO:17) adds a 3' DNA sequence for the joining region-kappa exon splice junction followed by a BstBI restriction site for cloning.

PCR reactions were performed with reagents as described above and with templates of 1-2 ng of PvuI (New England Biolabs) digested plasmid; each of these templates was denatured by an initial one minute incubation at 96°C. The heavy chain products were amplified by 35 thermal cycles of 55 or 60°C for 30 sec., 72°C for 30 sec., then 96°C for 1 min and a final extension step at 72°C for 5 min. The light chain products were amplified by 8 thermal cycles of 55 or 60°C for 30 sec., 72°C for 30 sec., then 96°C for 1 min followed by 30 step cycles of 60°C for 1 min, then 96°C for 1 min, and a final extension step at 72°C for 5 min.

The 13C4 heavy chain PCR product (approximately 400 bp) was purified using Qiaquick PCR Purification columns (Qiagen) as described by the manufacturer's instructions and subsequently digested with NruI and EcoRI (New England Biolabs). The digested PCR products were purified using the Wizard PCR Purification system (Promega) as per manufacturer's procedure and ligated into NruI /EcoRI digested and gel-purified pJRS3I3, resulting in plasmid tKMC229C (see Figure 4). The final consensus DNA sequence of the heavy chain variable region and proper splicing of the restriction sites were confirmed in this construct.

The 13C4 light chain PCR product (approximately 350 bp) was purified using Qiaquick PCR Purification columns (Qiagen) as described by the manufacturer's instructions and subsequently digested with EcoRV and BstBI (New England Biolabs). The digested PCR products were purified using Qiaquick PCR Purification columns (Qiagen) as per
5 manufacturer's procedure and ligated into EcoRV/BstBI digested and gel-purified tKMC180C2 (as described above), resulting in plasmid tKMC231D (see Figure 4). The final consensus DNA sequence of the light chain variable region and proper splicing of the restriction sites were confirmed in this construct.

EXAMPLE 3. Stable production of recombinant chimeric mouse/human 13C4 antibody.

10 A. Transfection of NSO Cells

The plasmid tKMC249A was transfected into NSO cells (Baxter International, Inc., Durante, CA) using electroporation after linearization with PvuI (New England Biolabs). 40 micrograms of the digested plasmid was mixed with 1×10^7 cells in a total volume of 800 microliters in a 0.4 centimeter cuvette and subjected to a pulse of 250mA, 960 μ F. The cells
15 were plated out after 24 hours into 96-well tissue culture plates, 6 plates with 200 μ l/well, and incubated at 37°C and 10% CO₂. As colonies appeared, the supernatants were assayed for the production of "humanized" antibody and for the capability of the expressed antibody to bind to Stx1.

B. Assay for Antibody Production

20 Antibody production and activity assays for the stable transfectants were performed as described below. These assays demonstrate that the transfection of cells with this plasmid construct can result in the production of a stable cell line that produces a humanized chimeric version of the 13C4 mouse hybridoma antibody (designated H13C4).

Antibody production ELISA assays were performed in 8-well strips from 96-well microtiter plates (Maxisorp F8; Nunc, Inc.) coated at a 1:500 dilution with Goat anti-Human IgG antibody (Pierce or Biodesign International) using a Tris-HCl coating buffer, pH 8.5. The plates were covered and incubated overnight at 4°C. Plates were then washed once with a wash storage buffer (Tris-HCl/NaCl/0.1% NaN₃). 50 microliters of culture supernatant was then applied to each well that had been filled previously with 50 microliters of a sample/conjugate diluent (Tris-HCl/NaCl/gelatin/Tween-20). The plates were allowed to incubate for 30 to 60 minutes on a rotator at room temperature. They were then washed three times with a wash solution (imidazole/NaCl/Tween-20). A goat anti-human kappa-HRP (Southern Biotechnologies) conjugate was diluted 1:250 in the sample/conjugate diluent and 100 microliters was added to the wells. The plates were incubated on a rotator for 30 to 60 minutes at room temperature. They were washed 6 times using the wash buffer, and then incubated with 100 µL/well of ABTS developing substrate (Kirkgaard & Perry Laboratories) for 8 minutes at room temperature. The reaction was stopped with 100 µL/well of diluted Quench buffer (Kirkgaard & Perry Laboratories). The absorbance value at 405 nm was determined using an automated microtiter plate ELISA reader (Ceres UV900HI, Bioteck, Winooski, Vermont). The controls for the ELISA assay were a human IgG1K myeloma protein (Biodesign International) and supernatant collected from non-transfected NSO cells. This assay (see Figure 8) demonstrates that the transfection of cells with this plasmid construct results in the cells producing a molecule containing both human heavy chain (IgG) and light chain (kappa) domains.

The supernatants were then assayed for the ability of the expressed antibodies to bind to Stx1 protein by ELISA. The activity assays were performed in 8-well strips from 96-well microtiter plates (Maxisorp F8; Nunc, Inc.) coated with approximately 0.1 µg/well purified

Stx1 received from Alison O'Brien (or obtained as described in Example 7). The plate coating and ELISA procedure was performed in the same manner as the antibody assay above with the substitution of TMB (Kirkgaard & Perry Laboratories) for ABTS as a developing substrate. The absorbance value at 450 nm was determined using an automated microtiter plate ELISA reader (Ceres UV900HI, Bioteck, Winooski, Vermont). As a positive control, the original mouse monoclonal antibody 13C4 was used, and assayed with a goat anti-mouse IgG conjugate (Jackson Laboratories) at a 1:2000 dilution. This assay (see Figure 9) demonstrates that the transfection of cells with this plasmid construct results in cells producing immunoglobulin that binds to Stx1. Neither mouse nor human IgG1K lacking the anti-Stx variable region bound the toxin.

Humanization of the EHEC Anti-Stx2 antibody: 11E10

The following examples 4-7 relate to anti-Stx2 and -Stx2 variant antibody 11E10 and its humanized counterpart H11E10.

EXAMPLE 4. Cloning of the 11E10 variable region cDNAs.

The hybridoma cell line producing the "11E10" antibody (Anti-Stx2) was deposited on August 1, 1990, at the American Type Culture Collection, Rockville, MD under Accession No. CRL 1987, and can be obtained from the ATCC, or, as here, from Dr. Alison O'Brien (for details of hybridoma preparation, see Perera, L.P. et al., *J. Clinical Microbiol.*, 26:2127-2131 (1988)). A vial of the cell line was thawed, washed with serum free medium and then resuspended in IMDM (Mediatech) complete media supplemented with 10% FBS (Irvine).

Total RNA was isolated from 1×10^7 "11E10" cells using the RNeasy RNA Isolation kit (Qiagen) following the manufacturer's procedure. The RNA was dissolved in 10 mM Tris,

0.1 mM EDTA (pH 8.4) containing 0.03 U/ μ g Prime RNase Inhibitor (5'-3') to a final concentration of 0.63 μ g/ μ l.

Figure 1 shows the strategy for cloning the variable region gene fragments and Figure 5 lists the oligonucleotide primers used. The "11E10" total RNA (2.5 μ g) was converted to cDNA by using Superscript II-MMLV Reverse Transcriptase (Life Technologies) according to manufacturer's procedures. The mouse light chain (JS153, JS154, SEQ ID NO:11 and 12) and mouse heavy chain (JS300, SEQ ID NO:6) were used as specific primers. The first strand cDNA synthesis products were then purified using a Centricon-30 concentrator device (Amicon). Of the 70 μ l of cDNA recovered, 3.5 μ l was used as template DNA for PCR.

10 Typical PCR amplification reactions (100 μ l) contained template DNA, 50 pmoles of the appropriate primers (JS153, JS154 and JS009, JS010, JS011, JS012, SEQ ID NO:7-12 for light chains, JS160, JS161, JS162 and JS001, JS002, JS003, JS004, JS008, SEQ ID NO:28-30, SEQ ID NO:2-5 for heavy chains), 2.5 units of ExTaq polymerase (PanVera), 1x ExTaq reaction buffer, 200 μ M dNTP, and 2 mM $MgCl_2$. The template was denatured by an initial

15 five minute incubation at 96°C. The products were amplified by 35 thermal cycles of 96°C for 1 min., 55°C for 30 sec., and 72°C for 30 sec, followed by 5 min. at 72°C. The PCR products from the successful reactions were purified using the Wizard PCR Purification system (Promega) as per manufacturer's procedure.

The heavy chain PCR products (approximately 400 bp) and the light chain PCR

20 products (approximately 350 bp) were then cloned into a bacterial vector for DNA sequence determination. Ligations of the PCR fragments were carried out into the pCR2.1 T/A style cloning vector (Invitrogen) following the manufacturer's procedures using a 3:1 insert to vector molar ratio. Two μ l of the ligation reactions were used to transform the INV α F' competent cells (Invitrogen) as per the manufacturer's procedure. Plasmid clones containing

DNA inserts were identified using diagnostic restriction enzyme digests, with EcoRI (New England Biolabs). The DNA sequence of plasmids containing the heavy chain inserts of the appropriate size (400 bp) was then determined. The final consensus DNA sequence of the heavy chain variable region of 11E10 is shown in Figure 6 with the CDRs indicated by

5 underlining.

The light chain plasmid clones needed to be further characterized because the hybridoma cell line that expresses the "11E10" antibody was made by fusing mouse splenocytes with SP20 myeloma cells. The SP20 cell line transcribes a pseudogene for the kappa light chain. The pseudogene transcript, when converted to cDNA by RT-PCR, contains

10 an AflIII restriction site. For this reason, the plasmid clones for the light chain variable region were digested with AflIII and those products that did not cut were then submitted for DNA sequencing. The final consensus sequence of the light chain variable region is shown in Figure 6, with the CDRs indicated by underlining.

The variable region gene fragments were re-amplified by PCR using primers that

15 adapted the fragments for cloning into the expression vector (see Figures 5 and 7). The heavy chain front primer (11E10HF, SEQ ID NO:37) includes a 5' tail that encodes the C-terminus of the heavy chain leader and an NruI restriction site for cloning, while the heavy chain reverse primer (11E10HB, SEQ ID NO:38) adds a 3' EcoRI restriction site for cloning. The light chain front primer (11E10LF, SEQ ID NO:39) includes a 5' tail that encodes the C-

20 terminus of the light chain leader and an EcoRV restriction site at the N-terminus of the light chain variable region for cloning, while the light chain reverse primer (11E10LB, SEQ ID NO:40) adds a 3' DNA sequence for the joining region-kappa exon splice junction followed by a BstBI restriction site for cloning. PCRs were performed as described above except,

following a 5 min. incubation at 96°C, the PCR parameters were 30 thermal cycles of 96°C for 1 min., 62°C for 30 sec., and 70°C for 30 sec., followed by 5 min. at 72°C.

The heavy chain variable region PCR product was then subcloned into a mammalian expression plasmid vector (pJRS315, produced as set forth below in Example 5) for production of recombinant chimeric mouse/human antibody molecules. The resulting vector clone was designated pACE1. Before the light chain variable region could be subcloned into the mammalian expression vector, it was subcloned into the pCR2.1 T/A style cloning vector previously described. The resulting plasmid was designated pACELC and digested, with BstBI/EcoRV, to cut out the light chain variable region. The variable region was then subcloned into the mammalian expression vector containing the "11E10" heavy chain variable region (pACE1). The final expression vector clone was designated pACE4. This vector results in the production of recombinant antibody molecules under the control of the CMV transcriptional promoters. The heavy chain molecules are direct cDNA constructs that fuse the variable region sequence directly into the human IgG1 constant domain. The light chain molecules, on the other hand, have a mouse kappa intron region 3' of the variable region coding fragment. After splicing, the variable region becomes fused to a human kappa constant region exon (see Figure 7). The selectable marker for the vector in mammalian cells is neomycin (G418).

For the production of pACE1, the "11E10" heavy chain PCR product (approximately 400 bp) was digested with NruI and EcoRI (New England Biolabs), purified using a Qiaquick PCR Purification column (Qiagen), as described by the manufacturer, and ligated into NruI/EcoRI digested and gel-purified pJRS315, resulting in plasmid pACE1 (see Figure 7).

The "11E10" light chain PCR product (approximately 350 bp) was cloned into the T/A cloning vector as per manufacturer's instructions. The resulting clone, pACELC, was

digested with EcoRV and BstBI (New England Biolabs) and the light chain fragment was gel-purified. This fragment was then ligated into the EcoRV/BstBI digested and gel-purified pACE1, resulting in plasmid pACE4 (see Figure 7). The sequence of the heavy and light chain variable regions was verified prior to mammalian cell transfection.

5 EXAMPLE 5. Construction of the expression vector pJRS315.

The plasmid pJRS315 is the expression plasmid into which the variable regions of the 11E10 antibody were cloned. This plasmid is a derivative of a basic expression vector, plasmid pSUN15, that contains no antibody variable region coding information, pJRS315 was created using DNA fragment ligations and site directed mutagenesis steps. The result was a
10 vector that expresses both antibody chains with CMV promoter driven transcription (see Figure 17). Neomycin resistance serves as the dominant selectable marker for transfection of mammalian cells. In addition, it has been designed to allow convenient cloning of any light chain variable region as an EcoRV/BstBI fragment, any heavy chain variable region as a NruI/EcoRI fragment, and any heavy chain constant domain as an EcoRI/NotI fragment.
15 These restriction sites were chosen because they occur rarely (if ever) in human and mouse variable regions. There is a mouse J region/kappa intron fragment fused to a human kappa exon so that after post-transcriptional splicing a mouse/human chimeric kappa light chain is produced.

The backbone of the vector was the plasmid pCDNA3 (Invitrogen). This plasmid was
20 cut with HindIII/XhoI and a "light chain polylinker" DNA fragment was inserted to create the starting "light chain vector". This linker contained the restriction sites HindIII, KpnI, ClaI, PmlI, EcoRV, XmaI, BamHI and XhoI to facilitate subsequent cloning steps to create the plasmid pCDNA3.LCPL. A SmaI/BclI DNA fragment containing a light chain leader, anti-

CKMB kappa light chain genomic fragment, and 3' UTR was cloned into the EcoRV/BamHI sites of pCDNA3.LCPL. The mouse kappa intron, exon and 3' UTR in this fragment was derived from LCPXK2 received from Dr. Richard Near (Near, R.I. et al., *Mol. Immunol.* 27:901-909, (1990)). Mutagenesis was then performed to eliminate an NruI (209), MluI 5 (229), and BstBI (2962) and to introduce an NheI (1229) and a BamHI (1214) site to create pcDNA3mut.LCPL.LCVK (see Figure 12).

A second "heavy chain vector" was constructed from the pcDNA3mut.LCPL.LCVK plasmid by replacing the light chain expression region (HindIII/XhoI) with a "heavy chain polylinker" consisting of restriction sites HpaI, BspEI, EcoRV, KpnI, and XhoI. This plasmid 10 was digested with EcoRV and KpnI. A SmaI/KpnI digested DNA fragment containing a heavy chain leader and an anti-CKMB IgG2b mouse heavy chain genomic fragment was then ligated into the EcoRV/KpnI digested plasmid. A KpnI/SalI oligonucleotide fragment containing a 3' UTR and a NotI upstream of the SalI site was subsequently cloned into the KpnI/XhoI digested plasmid (knocking out the XhoI site), to create the plasmid 15 pCDNA3mut.HCPL.HCV2b (see Figure 13).

From this point, two vectors were created that did not have any of the anti-CKMB variable or constant domain DNA sequences. This was done by cutting the plasmid pcDNA3mut.LCPL.LCVK with EcoRV/XhoI and inserting a linker oligonucleotide fragment containing EcoRV, BstBI, and XhoI sites to create pSUN9 (see Figure 14). In a similar way, 20 the anti-CKMB fragment in pCDNA3mut.HCPL.HCV2b (NruI/NotI) was replaced by a linker oligonucleotide fragment containing NruI, EcoRI and NotI sites to create pSUN10 (see Figure 15). A human kappa light chain constant domain was then cloned into pSUN9 as a BstBI/XhoI fragment, and a human IgG1 constant domain was cloned into pSUN10 as a EcoRI/NotI fragment.

A BglII/NheI fragment from the human heavy chain vector was then cloned into the human light chain vector cut with BamHI/NheI to create pSUN15 (see Figure 16).

The plasmid pJRS315 was then constructed using pSUN15 through the following process. A heavy chain variable region from another, unrelated, hybridoma cell line
5 (approximately 400 bp) was digested with NruI and EcoRI (New England Biolabs), purified using a Qiaquick PCR Purification column (Qiagen), as described by the manufacturer, and ligated into NruI /EcoRI digested and gel-purified pSUN15, resulting in plasmid pJRS311 (see Figure 16).

At this point, a BstBI/NotI (New England Biolabs) DNA fragment containing a mouse
10 kappa J-kappa intron fragment fused to a human kappa exon fragment was digested and gel-purified from the vector tKMC180C2. This fragment was ligated into the backbone of pJRS311 digested with BstBI/NotI and gel-purified resulting in the plasmid pJRS315 (see Figure 17).

EXAMPLE 6. Stable production of recombinant chimeric mouse/human "11E10" antibody.

15 A. Transfection of NSO Cells

The plasmid pACE4 was transfected into NSO cells using electroporation. The plasmid was linearized with a PvuI restriction enzyme digestion. 40 µg of digested plasmid DNA was mixed with 7×10^6 cells in a total volume of 400 µL and incubated at room temperature with gentle agitation for one minute. 10 µL of DMSO (Sigma) were added to a
20 final concentration of 1.25%. The cells/DNA/DMSO mix was transferred to a cold 0.4 centimeter cuvette and subjected to one pulse of 250 volts, 960 µF. The cells were transferred to one well of a six-well plate containing 5 ml of non-selective media supplemented with DMSO (final concentration of 1.25%). After 24 hours at 37°C and 10% CO₂, the cells were

plated out into 96-well microtiter plates. As colonies appeared, the supernatants were assayed for the production of "humanized" antibody and for the capability for the expressed antibody to bind to the Stx2 toxin.

B. Assay for Antibody Production

5 Antibody production and activity assays for the stable transfectants were performed in 8-well strips from 96-well microtiter plates (Maxisorp F8; Nunc, Inc.) coated with a 1:500 dilution of goat anti-Human F(ab')₂ anti-IgG antibody (Southern Biotechnology) using a bicarbonate coating buffer, pH 8.5. The plates were covered with pressure sensitive film (Falcon, Becton Dickinson) and incubated overnight at 4°C. Plates are then washed once
10 with wash solution (imidazole/NaCl/0.4% Tween-20). 100 µL of culture supernatant was then applied and allowed to incubate for 30 minutes on a plate rotator at room temperature. The plates were washed five times with wash solution (imidazole/NaCl/0.4% Tween-20). A goat anti-human kappa-HRP (Southern Biotechnology) conjugate was diluted 1:800 in the sample/conjugate diluent and 100 µL was added to the samples, then incubated on a plate
15 rotator for 30 minutes at room temperature. The samples were washed as above and then incubated with 100 µL per well of ABTS developing substrate (Kirkgaard & Perry Laboratories) and the absorbance value at 405 nm was determined using an automated microtiter plate ELISA Reader (Ceres UV900HI, BioTek Instruments, Winooski, Vermont). This assay (see Figure 10) demonstrates that the transfection of cells with this plasmid
20 construct results in cells producing a molecule containing both human IgG and kappa domains.

The supernatants were then assayed for the ability of the expressed antibodies to bind to EHEC Stx2 toxin. The activity assay was performed, as above, using plates coated at 1 µg/ml with Stx2 toxin (obtained as in Example 7 from Dr. O'Brien's lab) in a bicarbonate

coating buffer, pH 8.5. This assay demonstrates that the transfection of cells with this plasmid construct can result in the production of a humanized chimeric version of the 11E10 mouse monoclonal antibody which effectively binds Shiga toxin type 2 (Figure 11).

Together, these assays demonstrate that the transfection of cells with this plasmid
5 construct can result in the production of a stable cell line that produces a humanized chimeric version of the 11E10 mouse hybridoma antibody.

EXAMPLE 7. Verifying Biological and Immunological Efficacy of Humanized Antibodies to Shiga toxin.

A. Vero Cell Cytotoxicity Assay

10 The efficacy of the humanized antibodies to Shiga toxin obtained as detailed in Examples 1 through 6 was determined by assaying their ability to protect Vero cells from toxin. Cytotoxicity assays were performed essentially as described by Gentry and Dalrymple, *J. Clin. Microbiol.*, 12: 361-366 (1980). Briefly, toxin was obtained from cultures of *E. coli* K-12 strains that contained either plasmid pLPSH3 (encodes Stx; *J. Infect. Disease* 164:344-
15 352 (1991)) or pMJ 100 (encodes Stx2; *Inf. and Immunity*, 57:3743-3750 (1989)). Bacteria were disrupted by sonic lysis and clarified by centrifugation. The extracts were serially diluted in tissue culture medium (Dulbecco modified Eagle medium containing 10% fetal calf serum, 0.8 mM glutamine, 500 U of penicillin G per ml, and 500 mg of streptomycin per ml). One hundred microliters of 10-fold dilutions of the lysates were added to microtiter plate
20 wells containing about 10^4 Vero cells in 100 μ l of medium. The tissue culture cells were incubated at 37°C in 5% CO₂ for 48 hours and then fixed and stained with crystal violet. The intensity of color of the fixed and stained cells was measured with a Titertek reader at 620 nm. Incubation without antibody provides a standard toxicity curve for the Stx.

B. Antisera Neutralization Assay

Humanized mouse antibodies obtained according to the methods described in Examples 1 through 6 were tested for toxin neutralization. Neutralization of cytotoxic activity was described in great detail in Schmitt et al., *Infect. and Immun.*, 59:1065-1073

- 5 (1991). Briefly, lysates were incubated with serial dilutions of the humanized mouse antibodies at 37°C for 2 hours. One hundred microliters of the samples were then added to Vero cells as described above.

TABLE 1: Vero Cell Neutralization Assay

	Antibody	Required ng to Neut. 1 CD_{50} *
10	Mouse 13C4	25
	Humanized 13C4	26
	Mouse 11E10	2.7
	Humanized 11E10	82.8

*(1 CD_{50} = 1 pg pure toxin)

15 EXAMPLE 8. Passive Immunization with Humanized Antibodies

A. Protection against injection with crude Stx1 toxin.

At day zero, groups of five CD-1 mice were injected intraperitoneally (0.1 ml) with antibody H13C4 (humanized α -Stx1B), 13C4 (mouse α -Stx1B), H11E10 (humanized α -Stx2A), or phosphate buffered saline (PBS). These injections were repeated at day 1. The

20 mice were then challenged by intravenous injection with crude Stx1 toxin (obtained as described in Example 7) at doses of 1.7×10^5 or 1.7×10^6 CD_{50} . These doses of toxin were chosen following preliminary experiments with varying amounts of toxin. Mice were monitored for 21 days.

The results (Table 2) show clearly that the injected antibodies protect the mice against

25 at least 10 times the normal lethal toxin dose. As a negative control, the antibodies to Stx2

did not protect against Stx1 toxin challenge, indicating that the protection was specific to the toxin/antibody pair, and was not an artifact of the antibody preparation process.

TABLE 2: CD-1 Mice Injected Intravenously with Crude Stx1

	Antibody (dose/mouse)	#LD ₅₀ Protected Against	μg/kg of Ab to Protect from
			1LD ₅₀ *
5	Murine 13C4 (1.4 μg)	10	6.1
	Humanized 13C4 (4.1 μg)	20	8.9
	Humanized 11E10 (232 μg)	0	No protection

*(1 LD₅₀ = 30 ng crude Stx1)

B. Protection against oral infection with Stx2-producing EHEC strains.

10 Two different strains of mice and bacteria were used in these studies to test efficacy against both Stx2 and Stx2-variant. DBA/2J mice were challenged with EHEC strain 86-24 (O157:H7, Stx2⁺) and CD-1 mice were challenged with strain B2F1 (O91:H21, Stx2-variant⁻). While *E. coli* strain B2F1 is normally fatal to both mice strains, *E. coli* strain 86-24 is fatal to DBA/2J mice, while CD-1 mice will survive 86-24 infection.

15 At day zero, antibody H11E10 (humanized α-Stx2A) or 11E10 (mouse α-Stx2A) was injected intraperitoneally (0.1 ml) into groups of four or five mice. Control groups included mice that had received antibody 13C4 (mouse α-Stx1B), mice that had received 11E10 ascites fluid (mouse α-Stx2A), or mice that had received PBS instead of antibody. Mice were given streptomycin (5g/L) in their drinking water to decrease normal intestinal flora and their food
20 was removed. Streptomycin resistant derivatives of strains 86-24 (O157:H7, Stx2⁺) and B2F1 (O91:H21, Stx2-variant⁻) were grown overnight in L broth.

The following day (day 1) the mice received a second injection of test antibody, control antibody, or PBS. The mice were immediately fed 10¹⁰ CFU of 86-24 that had been pelleted and resuspended in 20% sucrose or 10³ CFU of B2F1 that had been serially diluted in

20% sucrose. Food was returned to the cage and the mice were monitored for 21 days.

(CFU=Colony Forming Units)

As shown in Table 3, immunization of the mice with either the murine or the humanized anti-Stx2 antibodies resulted in complete protection from a lethal oral dose of EHEC. Immunization with mouse 13C4 antibody, prepared in the same way but immunoreactive with the Stx1 toxin instead of the Stx2 toxin did not protect the mice from challenge with 86-24, a result that indicates the immunospecificity of the response.

TABLE 3: DBA/2J Mice Fed 10^{10} CFU 0157 (Stx2)

	Antibody (dose/mouse)	Survivors
10	Murine 13C4 (1.4 μ g)	0/5
	Murine 11E10 (1.0 μ g)	5/5
	Humanized 11E10 (1.0 μ g)	5/5

Similar results are illustrated in Table 4, where the more resistant CD-1 mice were fed the B2F1 *E. coli* strain which produces Stx2-variant toxin. Without treatment with antibody, mortality was total, but both the mouse and the humanized antibodies protected against Stx2-variant in a dose dependent manner.

TABLE 4: CD-1 Mice Fed 10^3 CFU O91:H21 (Stx2-variant)

	Antibody (dose/mouse)	Survivors
	PBS	0/5
20	Murine 11E10 (8.7 μ g)	1/4*
	Murine 11E10 (6.45 mg)	5/5
	Humanized 11E10 (23.2 μ g)	0/5
	Humanized 11E10 (232 μ g)	5/5

Protective dose between 10.1 and 1.0 mg/kg

*Delayed mean time to death by 3.4 days

EXAMPLE 9. Treatment of disease caused by bacteria producing Shiga toxin.

The subject invention also provides for a variety of methods of treating, ameliorating, or preventing, the diseases and effects associated with exposure to Shiga toxin. Positive clinical responses in humans have been obtained with monoclonal antibodies, and one skilled in the art would know how to employ anti-Stx humanized monoclonal antibodies in humans.

5 (See Fagerberg *et al.*, *Cancer Research*, 55:1824-27 (1995); *Eur. J. Cancer*, 2:261-267 (1995)). The precise dosage of humanized anti-Shiga toxin antibody administered to a patient for treatment of these diseases will vary in accordance with factors appreciated by the typical clinician. These factors include (but are not limited to) size, age, overall health, the extent of infection, and other medications being administered to said patient. The development of a

10 precise treatment regimen will require optimization through routine medical procedures well known to those in the medical arts. Examples of potential patient groups would include (but not be limited to) young children with bloody diarrhea but no white cells in the stool, patients with indications of HUS, patients with positive stool sample tests for Shiga toxin, siblings or daycare cohorts in contact with a case (as a passive preventative measure), and any patient

15 with diarrhea (not necessarily bloody) that has been in contact with an identified case. A typical dosage of about 5mg/kg body weight of humanized 13C4 combined with about 10mg/kg body weight of humanized 11E10 would be contemplated. This combined formulation could be administered to the patient twice to ensure effectiveness. Inclusion of both types of humanized antibodies together provides assurance that the patient will be

20 protected against all types of Shiga toxin.

The person skilled in the art would understand how to use and practice the invention based on the above disclosure. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as

exemplary only, with the true scope and spirit of the invention being indicated by the following claims.

We claim:

1. A humanized monoclonal antibody that binds to Shiga toxin protein, comprising a constant region and a variable region, wherein said constant region contains at least part of a human immunoglobulin constant region and said variable region contains at least part of a non-human immunoglobulin variable region.
2. The humanized monoclonal antibody of claim 1, having the same binding specificity as the antibody selected from the group consisting of murine 13C4 (ATCC Accession No. CRL 1794), murine 11E10 (ATCC Accession No. CRL 1987), humanized 13C4 (H13C4), and humanized 11E10 (H11E10).
3. The humanized monoclonal antibody of claim 1, wherein the antibody binds Shiga toxin type 1.
4. The humanized monoclonal antibody of claim 3, wherein said non-human variable region is from the mouse.
5. The humanized monoclonal antibody of claim 3, wherein at least part of said variable region is from the sequences as set forth in Figure 3 (SEQ ID NO:19 and SEQ ID NO:21).
6. A fragment of the antibody of claim 3, wherein the fragment binds Shiga toxin type 1.

7. The humanized monoclonal antibody of claim 3, wherein said human constant region is selected from the group consisting of IgG, IgA and IgM.

8. The humanized monoclonal antibody of claim 7, wherein said human constant region is IgG.

9. A humanized monoclonal antibody which binds Shiga toxin type 1, comprising a constant region and a variable region, wherein:

said constant region is IgG1-kappa, and

said variable region contains at least part of the sequence as set forth in Figure 3

(SEQ ID NO: 19 and SEQ ID NO: 21).

10. A humanized monoclonal antibody which binds Shiga toxin type 1, comprising a constant region and a variable region, wherein:

said constant region is IgG1-kappa, and

said variable region contains at least part of the CDR sequences as set forth in Figure

3,

said CDR sequences located as follows:

Heavy Chain CDRs:	CDR1-aa31-35
-------------------	--------------

(SEQ ID NO:19)	CDR2-aa50-66
----------------	--------------

	CDR3-aa99-111
--	---------------

Light Chain CDRs:	CDR1-aa24-34
(SEQ ID NO:21)	CDR2-aa50-56
	CDR3-aa89-97

11. An expression vector comprising a DNA sequence encoding the variable and constant regions of the light and the heavy chains of the antibody of claim 9, where the coding sequences for the variable regions contain at least part of the DNA sequences as set forth in Figure 3 (SEQ ID NO:18 and SEQ ID NO:20).

12. A host cell transformed with the expression vector of claim 11.

13. The humanized monoclonal antibody of claim 1, wherein the antibody binds Shiga toxin type 2 and Shiga toxin type 2 variants.

14. The humanized monoclonal antibody of claim 13, wherein said non-human variable region is from the mouse.

15. The humanized monoclonal antibody of claim 13, wherein at least part of said variable region is from the sequence as set forth in Figure 6 (SEQ ID NO: 42 and SEQ ID NO:44).

16. A fragment of the antibody of claim 13, wherein the fragment binds Shiga toxin type 2 and Shiga toxin type 2 variants.

17. The humanized monoclonal antibody of claim 13, wherein said human constant region is selected from group consisting of IgG, IgA and IgM.

18. The humanized monoclonal antibody of claim 16, wherein said human constant region is IgG.

19. A humanized monoclonal antibody which binds Shiga toxin type 2 and Shiga toxin type 2 variants, comprising a constant region and a variable region, wherein:

said constant region is IgG1-kappa, and

said variable region contains at least part of the sequence as set forth in Figure 6

(SEQ ID NO:42 and SEQ ID NO:44).

20. A humanized monoclonal antibody which binds Shiga toxin type 2 and Shiga toxin type 2 variants, comprising a constant region and a variable region, wherein:

said constant region is IgG1-kappa, and

said variable region contains at least part of the CDR sequences as set forth in Figure

6,

said CDR sequences located as follows:

Heavy Chain CDRs: CDR1-aa31-35

(SEQ ID NO:44) CDR2-aa50-66


CDR3-aa99-108

Light Chain CDRs:	CDR1-aa24-40
(SEQ ID NO:42)	CDR2-aa56-62
	CDR3-aa95-103

21. An expression vector comprising a DNA encoding the variable and constant regions of the light and the heavy chains of the antibody of claim 19, where the coding sequences for the variable regions contain at least part of the DNA sequences as set forth in Figure 6 (SEQ ID NO:41 and 43).
22. A host cell transformed with the expression vector of claim 20.
23. A pharmaceutical composition comprising the antibody of claim 1, or fragment or derivative thereof, and a pharmaceutically acceptable carrier or diluent.
24. A method for treating a patient having an infection caused by EHEC or other Shiga toxin producing bacteria comprising administering to the patient a therapeutically effective amount of the pharmaceutical composition of claim 22.
25. A method for reducing illness caused by EHEC or other Shiga toxin-producing bacteria comprising administering to the patient a prophylactically effective amount of the pharmaceutical composition of claim 22.
26. A pharmaceutical composition comprising the antibody of claim 3, or fragment or derivative thereof, and a pharmaceutically acceptable carrier or diluent.

27. A method for treating a patient having an infection caused by EHEC or other Shiga toxin producing bacteria comprising administering to the patient a therapeutically effective amount of the pharmaceutical composition of claim 25.
28. A method for reducing illness caused by EHEC or other Shiga toxin-producing bacteria comprising administering to the patient a prophylactically effective amount of the pharmaceutical composition of claim 25.
29. A pharmaceutical composition comprising the antibody of claim 13, or fragment or derivative thereof, and a pharmaceutically acceptable carrier or diluent.
30. A method for treating a patient having an infection caused by EHEC or other Shiga toxin producing bacteria comprising administering to the patient a therapeutically effective amount of the pharmaceutical composition of claim 28.
31. A method for reducing illness caused by EHEC or other Shiga toxin-producing bacteria comprising administering to the patient a prophylactically effective amount of the pharmaceutical composition of claim 28.

1/18



2/18

MOUSE HEAVY CHAIN "FRONT" PRIMERS

JSS1

5'-ATTTCAGGCCCAGCCGGCCATGGCCGARGTRMAGCTKSAKGAGWC-3' SEQ ID NO.1

JSS2

5'-ATTTCAGGCCCAGCCGGCCATGGCCGARGTYCARCTKCARCARYC-3' SEQ ID NO.2

JSS3

5'-ATTTCAGGCCCAGCCGGCCATGGCCAGGTGAAGCTKSTSGARTC-3' SEQ ID NO.3

JSS4

5'-ATTTCAGGCCCAGCCGGCCATGGCCGAVGTGMWGCTKGTGGAGWC-3' SEQ ID NO.4

JSS8

5'-ATTTCAGGCCCAGCCGGCCATGGCCAGGTBCARCTKMARSARTC-3' SEQ ID NO.5

MOUSE HEAVY CHAIN "BACK" PRIMERS

JS300

5'-GAARTAVCCCTTGACCAGGC-3' SEQ ID NO.6

MOUSE LIGHT CHAIN LEADER "FRONT" PRIMERS

JSS009

5'-GGAGGCGGCGGTTCTGACATTGTGMTGWCMCARTC-3' SEQ ID NO.7

JSS010

5'-GGAGGCGGCGGTTCTGATRRTTKYGATGACBCARRC-3' SEQ ID NO.8

JSS011

5'-GGAGGCGGCGGTTCTGAYATYMAGATGACMCAGWC-3' SEQ ID NO.9

JSS012

5'-GGAGGCGGCGGTTCTSA AATTGWKCTSACYCAGTC-3' SEQ ID NO.10

FIG. 2A

3/18

MOUSE LIGHT CHAIN "BACK" PRIMER

JS153

5'-TTCATAGGCGGCCGCACTAGTAGCMCGTTTCAGYTCCARC-3'

SEQ ID NO.11

JS154

5'-TTCATAGGCGGCCGCACTAGTAGCMCGTTTKATYTCCARC-3'

SEQ ID NO.12

OKA57

5'-GCACCTCCAGATGTAACTGCTC-3'

SEQ ID NO.13

"13C4" SPECIFIC PRIMERS

OKA144

5'-CTTGATCGCGACAGCTACAGGTGTCCACTCCCAGGTGCAGCTGCAGGAG-3' SEQ ID NO.14

OKA143

5'-GGTATGGAATTCTGAGGAGACTGTGAGAGTGGTGCC-3'

SEQ ID NO.15

OKA145

5'-GGTTCTGATATCGTGATGTCCCAGTCTCACAAATTC-3'

SEQ ID NO.16

OKA146

5'-GACATATTCGAAAAGTGTACTTACGTTTCAGCTCCAGACTGG-3'

SEQ ID NO.17

FIG. 2B

4/18

13C4 ANTI-STX1 HEAVY CHAIN VARIABLE REGION


CAGGTGCAGCTGCAGGAGTCTGGGGCTGAGCTGGTGAGGTCTGGGGCCTCAGTGAGGATG SEQ ID NO.18
Q V Q L Q E S G A E L V R S G A S V R M SEQ ID NO.19
TCCTGCAAGGCTTCTGGCTACACATTTACCAGTTACAATATGCACTGGGTAAAACAGACA
S C D A S G Y T F T S Y N M H W V K Q T
CCTGGACAGGGCCTGGAATGGATTGGATATATTTATCCTGGAAATGGTGGTACTAACTAC
P G Q G L E W I G Y I Y P G N G G T N Y
ATTCAGAAATTTAAGGGCAAGGCCATATTGACTGCAGACACATCCTCCAGCACAGCCTAC
I Q K F K G K A I L T A D T S S S T A Y
ATGCAGATCAGCAGTCTGACATCTGAAGACTCTGCGGTCTATTTCTGTACAAGAAGTCCTCT
M Q I S S L T S E D S A V Y F C T R S P S
CACTACAGTAGTGACCCCTACTTTGACTACTGGGGCCAGGGCACCCTCTCACAGTCTCCTCA
H Y S S D P Y F D Y W G Q G T T L T V S S

13C4 ANTI-STX1 LIGHT CHAIN VARIABLE REGION

GATATCGTGATGTCCCAGTCTCACAAATTCATGTCCACATCAGTCGGAGACAGGGTCAGCATC SEQ ID NO.20
D I V M S Q S H K F M S T S V G D R V S I SEQ ID NO.21
ACCTGTAAGGCCAGCCAGGATGTGGGTACTGCTGTTGCCTGGTATCAGCAGAATCCAGGACAA
T C K A S Q D V G T A V A W Y Q Q N P G Q
TCTCCTAAATTTCTGATTTACTGGGCATCCACACGGCACACTGGAGTCCCTGATCGCTTCACA
S P K F L I Y W A S T R H T G V P D R F T
GGCAGTGGATCTGGGACAGATTTCACTCTCACCATTACCAATGTGCAGTCTGAAGACTTGGCA
G S G S G T D F T L T I T N V Q S E D L A
GATTATTTCTGTGAGCAATATAGCAGTTATCCTCTCACGTTTCGGTGTGGGACCAGTCTGGAG
D Y F C Q Q Y S S Y P L T F G A G T S L E
CTGAAACGT
L K R

FIG. 3

5/18

**FIG. 4**

6/18

MOUSE HEAVY CHAIN "FRONT" PRIMERS

JS001

5'-ATTTTCAGGCCCCAGCCGGCCATGGCCGARGTRMAGCTKSAKGAGWC-3' SEQ ID NO.22

JS002

5'-ATTTTCAGGCCCCAGCCGGCCATGGCCGARGTYCARCTKCARCARYC-3' SEQ ID NO.23

JS003

5'-ATTTTCAGGCCCCAGCCGGCCATGGCCGAGGTGAAGCTKSTSGARTC-3' SEQ ID NO.24

JS004

5'-ATTTTCAGGCCCCAGCCGGCCATGGCCGAVGTGMWGCTKGTGGAGWC-3' SEQ ID NO.25

JA008

5'-ATTTTCAGGCCCCAGCCGGCCATGGCCGAGGTBCARCTKMARSARTC-3' SEQ ID NO.26

MOUSE HEAVY CHAIN "BACK" PRIMERS

JS300 5'-GAARTAVCCCTTGACCAGGC-3' SEQ ID NO.27

JS160 5'-GCTGCCACCGCCACCTGMRGAGACDGTGASTGARG-3' SEQ ID NO.28

JS161 5'-GCTGCCACCGCCACCTGMRGAGACDGTGASMTRG-3' SEQ ID NO.29

JS162 5'-GCTGCCACCGCCACCTGMRGAGACDGTGASCAGRG-3' SEQ ID NO.30

MOUSE LIGHT CHAIN LEADER "FRONT" PRIMERS

JS009 5'-GGAGGCGGCGGTTCTGACATTGTGMTGWCMCARTC-3' SEQ ID NO.31

JS010 5'-GGAGGCGGCGGTTCTGATRTTKYGATGACBCARRC-3' SEQ ID NO.32

JS011 5'-GGAGGCGGCGGTTCTGAYATYMGATGACMCAGWC-3' SEQ ID NO.33

JS012 5'-GGAGGCGGCGGTTCTSAATTGWKCTSACYCAGTC-3' SEQ ID NO.34

FIG. 5A

7/18

MOUSE LIGHT CHAIN "BACK" PRIMERS

JS153 5'-TTCATAGGCGGCCGCACTAGTAGCMCGTTTCAGYTCCARC-3' SEQ ID NO.35

JS154 5'-TTCATAGGCGGCCGCACTAGTAGCMCGTTTKATYTCCARC-3' SEQ ID NO.36

"11E10" SPECIFIC PRIMERS

11E10HF SEQ ID NO.37

5'-ATATACTCGCGACAGCTACAGGTGTCCACTCCGAAGTCCAACGCAACAGCC-3'

11E10HB

5'-ATTAATGAATTCTCGGAGACGGTGAGAGTGGTC-3' SEQ ID NO.38

11E10LF

5'-TTAAATGATATCGTGCTGTCACAATCTCC-3' SEQ ID NO.39

11E10LB

5'-TAATCGTTCGAAAAGTGTACTTACGTTTCAGTTCAGCTTGGTCC-3' SEQ ID NO.40

FIG. 5B

8/18

11E10 LIGHT CHAIN 7/23/97 12:38 PM										DOUBLE UNDERLINED = CDRs										SEQ ID NO. 41-42	
		10	20	30	40	50	60	70	80												
21	11E10	GACATTGTGC TGTCACAATC TCCATCCTCC CTAGTTGTGT CAGTTGGAGA GAAGTTACT ATGAGCTGCA AGTCTAGTCA																			
27		D I V L S Q S P S S L V V S V G E K V T M S C K S S Q																			
		90	100	110	120	130	140	150	160												
21	11E10	GAGCCTTTTA TATAGTAGAA ATCAAAGAA CTACTTGCC TGGTACCAGC AGAAACCAGG GCAGTCTCCT AAAGTGCTGA																			
27		S L L Y S R N Q K N Y L A W Y Q Q K P G Q S R K V L																			
		170	180	190	200	210	220	230	240												
21	11E10	TTTACTGGC ATCTACTAGG GAATCTGGG TCCCTGATCG CCTCACAGC AGTGGATCTG GGACAGATTT CACTCTCACC																			
27		I Y W A S T R E S G V P D R L T G S G S G T D F T L T																			
		250	260	270	280	290	300	310	320												
21	11E10	ATCAGCAGTG TGAAGGCTGA AGACCTGGCA GTTTATTACT GTCAGCAATA TTATAGTTAT CCGTCACGT TCGGTGCTGG																			
27		I S S V K A E D L A V Y Y C Q Q Y Y S Y P L T F G A G																			
		330	340	350	360	370	380	390	400												
21	11E10	GACCAAGCTG GAGCTGAAA																			
27		T K L E L K																			

FIG. 6A

9/18

11E10 HEAVY CHAIN 7/23/97 12:36 PM		DOUBLE UNDERLINED = CDRs										SEQ ID NO.43-44
		10	20	30	40	50	60	70	80			
1 11E10	GAAGTCCAAC TGCACACAGCC TGGACCTGAG CTGGAGAAGC CTGGCGCTTC AGTGAAGTTC TCCTGCAAGG CTTCGTGTTA											
11	E V Q L Q Q P G P E L E K P G A S V K L S C K A S G Y											
		90	100	110	120	130	140	150	160			
1 11E10	CTCTTTCACT GACTACAACA TGAACCTGGT GAAGCAGAAC AATGGAGAGA GCCTTGAGTG GATTGGAAAA ATTGATCCTT											
11	S F T D Y N M N W V K Q N N G E S L E W I G K I D P											
		170	180	190	200	210	220	230	240			
1 11E10	ACTATGGTGG TCCTAGCTAC AACCAGAAGT TCAAGGACAA GGCCACATTG ACTGTAGACA AGTCTTCCAG CACAGCCAC											
11	Y Y G G P S Y N Q K F K D K A T L T V D K S S S T A Y											
		250	260	270	280	290	300	310	320			
1 11E10	ATGCAGTTCA AGAGCTGAC ATCTGAGGAC TCTGCAGTCT ATTACTGTAC AAGAGGGGGA AATAGGGACT GGTACTTCGA											
11	M Q F K S L T S E D S A V Y Y C T R G G N R D W Y F D											
		330	340	350	360	370	380	390	400			
1 11E10	TGTGTGGGC GCAGGGACCA CGCTCACCGT CTCCGCA											

FIG. 6B

10/18




FIG. 7

11/18





FIG. 8

12/18

ELISA FOR STX1

**FIG. 9**

13/18




FIG. 10

14/18






FIG. 11

15/18

pCDNA3mut.LCPL.LCVK
construction:


**FIG. 12**

pCDNA3mut.HCPL.HCV2b
construction:


**FIG. 13**

16/18

pSUN9, 9:Kappa and pSUN10, 10:IgG1
constructions:


**FIG. 14****FIG. 15**

17/18

**FIG. 16**

SUBSTITUTE SHEET (RULE 26)

18/18

pJRS315
construction:**FIG. 17**

SEQUENCE LISTING

<110> HENRY JACKSON FOUNDATION FOR THE ADVANCEMENT OF MILITARY
MEDICINE

<120> HUMANIZED MONOCLONAL ANTIBODIES THAT PROTECT AGAINST
SHIGA TOXIN INDUCED DISEASE

<130> 04995.0032-00304

<140>

<141>

<160> 44

<170> PatentIn Ver. 2.0

<210> 1

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 1

att tca ggc cca gcc ggc cat ggc cga rgt rma gct ksa kga gwc 45

<210> 2

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 2

atttcaggcc cagccggcca tggccgargt ycarctkcar caryc 45

<210> 3
<211> 45
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide
<400> 3
atttcaggcc cagccggcca tggcccaggt gaagctksts gartc 45

<210> 4
<211> 45
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide
<400> 4
atttcaggcc cagccggcca tggccgargt rmagctksak gagwc 45

<210> 5
<211> 45
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide
<400> 5
atttcaggcc cagccggcca tggcccaggt bcarctkmar sartc 45

<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide
<400> 6
gaartavccc ttgaccaggc 20

<210> 7
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 7
ggaggcggcg gttctgacat tgtgmtgwcw cartc

35

<210> 8
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 8
ggaggcggcg gttctgatrt tkygatgacb carrc

35

<210> 9
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 9
ggaggcggcg gttctgayat ymagatgacw cagwc

35

<210> 10
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 10
ggaggcggcg gttctsaaat tgwktsacy cagtc

35

<210> 11
<211> 40
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 11
ttcataggcg gccgcactag tagcmcgttt cagytccarc

40

<210> 12
<211> 40
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 12
ttcataggcg gccgcactag tagcmcgttt katytccarc

40

<210> 13
<211> 23
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 13
gcacctccag atgttaactg ctc

23

<210> 14
<211> 49
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 14
cttgatcgcg acagctacag gtgtccactc ccagggtgcag ctgcaggag

49

<210> 15
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 15
ggtatggaat tctgaggaga ctgtgagagt ggtgcc 36

<210> 16
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 16
ggttctgata tcgtgatgtc ccagtctcac aaattc 36

<210> 17
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 17
gacatattcg aaaagtgtac ttacgtttca gctccagact gg 42

<210> 18
<211> 366
<212> DNA
<213> Shigella dysenteriae

<400> 18
caggtgcagc tgcaggagtc tggggctgag ctggtgaggt ctggggcctc agtgaggatg 60
tcctgcaagg cttctggcta cacatttacc agttacaata tgcactgggt aaaacagaca 120

cctggacagg gcctggaatg gattggatat atttatcctg gaaatggtgg tactaactac 180
 attcagaaat ttaagggcaa ggccatattg actgcagaca catcctccag cacagcctac 240
 atgcagatca gcagtctgac atctgaagac tctgcggtct atttctgtac aagaagtccc 300
 tctcactaca gtagtgaccc ctactttgac tactggggcc agggcaccac tctcacagtc 360
 tcctca 366

<210> 19

<211> 122

<212> PRT

<213> Shigella dysenteriae

<400> 19

Gln Val Gln Leu Gln Glu Ser Gly Ala Glu Leu Val Arg Ser Gly Ala

1 5 10 15

Ser Val Arg Met Ser Cys Asp Ala Ser Gly Tyr Thr Phe Thr Ser Tyr

20 25 30

Asn Met His Trp Val Lys Gln Thr Pro Gly Gln Gly Leu Glu Trp Ile

35 40 45

Gly Tyr Ile Tyr Pro Gly Asn Gly Gly Thr Asn Tyr Ile Gln Lys Phe

50 55 60

Lys Gly Lys Ala Ile Leu Thr Ala Asp Thr Ser Ser Ser Thr Ala Tyr

65 70 75 80

Met Gln Ile Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys

85 90 95

Thr Arg Ser Pro Ser His Tyr Ser Ser Asp Pro Tyr Phe Asp Tyr Trp

100 105 110

Gly Gln Gly Thr Thr Leu Thr Val Ser Ser

115 120

<210> 20
 <211> 324
 <212> DNA
 <213> Shigella dysenteriae

<400> 20
 gatatcgtga tgtcccagtc tcacaaattc atgtccacat cagtcggaga cagggtcagc 60
 atcacctgta aggccagcca ggatgtgggt actgctgttg cctgggtatca gcagaatcca 120
 ggacaatctc ctaaatttct gatttactgg gcatccacac ggcacactgg agtccttgat 180
 cgcttcacag gcagtggatc tgggacagat ttcactctca ccattaccaa tgtgcagtct 240
 gaagacttgg cagattatct ctgtcagcaa tatagcagtt atcctctcac gttcgggtgct 300
 gggaccagtc tggagctgaa acgt 324

<210> 21
 <211> 108
 <212> PRT
 <213> Shigella dysenteriae

<400> 21
 Asp Ile Val Met Ser Gln Ser His Lys Phe Met Ser Thr Ser Val Gly
 1 5 10 15
 Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Gly Thr Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Asn Pro Gly Gln Ser Pro Lys Phe Leu Ile
 35 40 45
 Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Thr Asn Val Gln Ser
 65 70 75 80
 Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Ser Ser Tyr Pro Leu
 85 90 95
 Thr Phe Gly Ala Gly Thr Ser Leu Glu Leu Lys Arg
 100 105

<210> 22
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 22
atttcaggcc cagccggcca tggccgargt rmagctksak gagwc 45

<210> 23
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 23
atttcaggcc cagccggcca tggccgargt ycarctkcar caryc 45

<210> 24
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 24
atttcaggcc cagccggcca tggcccaggt gaagctksts gartc 45

<210> 25
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 25
atttcaggcc cagccggcca tggccgavgt gmwgctkgtg gagwc 45

<210> 26
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 26
atttcaggcc cagccggcca tggcccaggt bcarctkmar sartc 45

<210> 27
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 27
gaartavccc ttgaccaggc 20

<210> 28
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 28
gctgccaccg ccacctgmrg agacdgtgas tgarg 35

<210> 29
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 29
gctgccaccg ccacctgmrg agacdgtgas mgtrg 35

<210> 30
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 30
gctgccaccg ccacctgmrg agacdgtgas cagrg 35

<210> 31
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 31
ggaggcggcg gttctgacat tgtgmtgwm cartc 35

<210> 32
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 32
ggaggcggcg gttctgatrt tkygatgacb carrc 35

<210> 33
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 33
ggaggcggcg gttctgayat ymagatgacm cagwc 35

<210> 34
<211> 35
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 34
ggaggcggcg gttctsaaat tgwktsacy cagtc

35

<210> 35
<211> 40
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 35
ttcataggcg gccgcactag tagcmggtt cagytccarc

40

<210> 36
<211> 40
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 36
ttcataggcg gccgcactag tagcmggtt katytccarc

40

<210> 37
<211> 52
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 37
atatactcgc gacagctaca ggtgtccact ccgaagtcca actgcaacag cc

52

<210> 38
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 38
attaatgaat tctgcggaga cggtagagat ggtc 34

<210> 39
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 39
ttaaatgata tcgtgctgtc acaatctcc 29

<210> 40
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 40
taatcggtcg aaaagtgtac ttacgtttca gttccagctt ggtcc 45

<210> 41
<211> 339
<212> DNA
<213> Shigella dysenteriae

<400> 41
gacattgtgc tgtcacaatc tccatcctcc ctagtttgtt cagttggaga gaaggttact 60
atgagctgca agtctagtca gaggctttta tatagtagaa atcaaaagaa ctacttggcc 120
tggtaccagc agaaaccagg gcagtctcct aaagtgtgta ttactgggc atctactagg 180

gaatctgggg tccctgatcg cctcacaggc agtggatctg ggacagattt cactctcacc 240
 atcagcagtg tgaaggctga agacctggca gtttattact gtcagcaata ttatagttat 300
 ccgctcacgt tcggtgctgg gaccaagctg gagctgaaa 339

<210> 42

<211> 113

<212> PRT

<213> Shigella dysenteriae

<400> 42

Asp Ile Val Leu Ser Gln Ser Pro Ser Ser Leu Val Val Ser Val Gly
 1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

Arg Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

Ser Pro Lys Val Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60

Pro Asp Arg Leu Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
 85 90 95

Tyr Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
 100 105 110

Lys

<210> 43

<211> 357

<212> DNA

<213> Shigella dysenteriae

<400> 43

gaagtccaac tgcaacagcc tggacctgag ctggagaagc ctggcgcttc agtgaaacta 60

tcctgcaagg cttctgggta ctctttcact gactacaaca tgaactgggt gaagcagaac 120

aatggagaga gccttgagtg gattggaaaa attgatacctt actatggtgg tcctagctac 180
 aaccagaagt tcaaggacaa ggccacattg actgtagaca agtcttccag cacagcctac 240
 atgcagttca agagcctgac atctgaggac tctgcagtct attactgtac aagaggggga 300
 aatagggact ggtacttcca tgtgtggggc gcagggaacca cgctcaccgt ctccgca 357

<210> 44

<211> 119

<212> PRT

<213> Shigella dysenteriae

<400> 44

Glu Val Gln Leu Gln Gln Pro Gly Pro Glu Leu Glu Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
 20 25 30

Asn Met Asn Trp Val Lys Gln Asn Asn Gly Glu Ser Leu Glu Trp Ile
 35 40 45

Gly Lys Ile Asp Pro Tyr Tyr Gly Gly Pro Ser Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Phe Lys Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Thr Arg Gly Gly Asn Arg Asp Trp Tyr Phe Asp Val Trp Gly Ala Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ala
 115

INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/US 98/27267

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/85 C07K16/12 C07K16/46 C12N5/10 A61K39/40		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	H. RÜSSMANN ET AL.: "Variants of Shiga-like toxin II constitute a major toxin component in Escherichia coli 0157 strains from patients with haemolytic uraemic syndrome." JOURNAL OF MEDICAL MICROBIOLOGY, vol. 40, no. 5, May 1994, pages 338-343, XP002100300 London, GB see the whole document --- -/--	1-31
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 19 April 1999		Date of mailing of the international search report 06/05/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Nooij, F

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/US 98/27267

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	A. LOBUGLIO ET AL.: "Mouse/human chimeric monoclonal antibody in man: Kinetics and immune response." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE U.S.A., vol. 86, June 1989, pages 4220-4224, XP002100301 Washington, DC, USA cited in the application see abstract	1-31
A	--- J. SPEIRS ET AL.: "Detection of Escherichia coli cytotoxins by enzyme-linked immunosorbent assays." CANADIAN JOURNAL OF MICROBIOLOGY, vol. 37, no. 8, August 1991, pages 650-653, XP002100302 Ottawa, Canada see abstract	1-31
A	--- EP 0 239 400 A (G.WINTER) 30 September 1987 see examples see claims	1-31
P,X	--- WO 98 20903 A (TRUSTEES OF TUFTS COLLEGE) 22 May 1998 see examples see claims -----	1,3,4, 6-8,13, 14, 16-18, 23-31

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/27267

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 24,25,27,28,30,31
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 24,25,27,28,30,31
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/US 98/27267

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 239400	A	30-09-1987	AT 109508 T	15-08-1994
			DE 3750304 D	08-09-1994
			DE 3750304 T	22-12-1994
			ES 2056814 T	16-10-1994
			GB 2188638 A,B	07-10-1987
			JP 62296890 A	24-12-1987
			US 5225539 A	06-07-1993
<hr/>				
WO 9820903	A	22-05-1998	NONE	
<hr/>				

This Page Blank (uspto)